

RESEARCH

Open Access



One-year outcomes in sepsis: a prospective multicenter cohort study in Japan

Keibun Liu^{1,32*†}, Shinichi Watanabe^{2,3†}, Kensuke Nakamura^{4,5}, Hidehiko Nakano⁵, Maiko Motoki⁵, Hiroshi Kamijo⁶, Matsuoka Ayaka⁷, Kenzo Ishii⁸, Yasunari Morita⁹, Takashi Hongo^{10,11}, Nobutake Shimojo¹², Yukiko Tanaka¹³, Manabu Hanazawa¹⁴, Tomohiro Hamagami¹⁵, Kenji Oike¹⁶, Daisuke Kasugai¹⁷, Yutaka Sakuda¹⁸, Yuhei Irie¹⁹, Masakazu Nitta²⁰, Kazuki Akieda²¹, Daigo Shimakura²², Hajime Katsukawa²³, Toru Kotani²⁴, David McWilliams^{25,26}, Peter Nydahl^{27,28}, Stefan J. Schaller^{29,30}, Takayuki Ogura³¹ and the ILOSS Study Group

Abstract

Background Sepsis is a leading cause of death in intensive care units (ICU). Sepsis survivors are often left with significant morbidity, termed post-intensive care syndrome (PICS), impacting post-sepsis life. The aim was to present detailed data on the prognostic and functional long-term outcomes of ICU patients with sepsis in Japan, which is currently lacking and therefore prevents development of targeted solutions.

Methods A multicenter prospective study, involving 21 ICUs in 20 tertiary hospitals in Japan, included all consecutive adult ICU patients between November 2020 and April 2022, and diagnosed with sepsis at ICU admission (Sepsis 3). Follow-ups were performed at 3, 6, and 12 months after hospital discharge by telephone and mail. Primary outcome was death or incidence of PICS, defined by any of physical dysfunction (Barthel Index ≤ 90), cognitive dysfunction (Short Memory Questionnaire < 40), or mental disorder (any subscales for anxiety or depression of Hospital Anxiety and Depression Scale ≥ 8 , or Impact of Event Scale-Revised ≥ 25). Secondary outcomes included Quality of Life (QOL), employment, and use of hospital, emergency, rehabilitation, and psychiatric services. A multivariable analysis investigated independent factors associated with each dysfunction at each follow-up.

Results A total of 339 patients were included (median age 74 [67–82] years, 60% male, 77% septic shock, and a median SOFA of 9 [6–12]). Mortality was 23% at hospital discharge, increasing to 37% at 12 months. The rate of death for those who met PICS Criteria at hospital discharge was 89%, with a death or PICS incidence of 73%, 64%, and 65% at 3, 6, and 12 months, respectively. Limited improvements in QOL and return to work (44%), high rates of hospital readmissions (40%), frequent emergency service usage (31%), and low utilization of rehabilitation and psychiatric services (15% and 7%) were identified over the first year. The incidence of any PICS-related dysfunction was consistently an independent factor for the incidence of the same dysfunction at the following follow-ups.

Conclusions This multicenter study identified the distinct realities of post-sepsis life in Japanese ICU patients, highlighting the unique challenges in improving their functions and returning to daily life.

Trial Registration University Hospital Medical Information Network UMIN000041433

Keywords Intensive care unit, Morbidity, Mortality, Post-intensive care syndrome, Quality of life, Sepsis

[†]Keibun Liu and Shinichi Watanabe equally contributed to this paper.

*Correspondence:

Keibun Liu

keiliu0406@gmail.com

Full list of author information is available at the end of the article



© The Author(s) 2025. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Background

Sepsis is a life-threatening condition characterized by an excessive immune response and subsequent organ dysfunction, resulting in high mortality and morbidity [1, 2]. Since 2020, a total of 48.9 million people worldwide have been affected, leading to 11 million sepsis-related deaths, which account for 20% of all global fatalities [3]. Due to advancements in sepsis management, survival rates have improved over the past few decades [4]. However, the morbidity experienced by sepsis survivors has been increasingly recognized, the impact of which extends beyond hospitalization and persists even after discharge, severely affecting essential life functions [5]. This long-term perturbation is known as Post-Intensive Care Syndrome (PICS), which is characterized by physical and cognitive impairments as well as mental disorders that can last for years following recovery from the Intensive Care Unit (ICU) [6–8]. PICS imposes a significant burden on the lives of patients post-sepsis, hindering their ability to return to their previous lives and jobs because of impaired activities of daily living (ADL) and a diminished quality of life (QOL) [9, 10]. Given the increasing number of sepsis patients, along with the decreasing trend of mortality in sepsis [11], effective interventions to prevent or treat sepsis-related morbidity are urgently required.

The number of studies exploring post-sepsis recovery following ICU and hospital discharge has significantly increased, primarily from Western countries outside of Japan. The characteristics and backgrounds of ICU admissions vary widely by region. For instance, the ICU population reported in Western contexts tends to have a relatively high BMI and younger age compared to the typically older age and lower BMI of patients in Japan's ICUs [12–14]. This heterogeneity in patient profiles may influence overall outcomes, posing challenges in interpreting studies from outside Japan that may not fully apply to the ICU population there. For example, multiple international studies have indicated that older ICU patients suffering from sepsis experienced significantly higher rates of mortality and morbidity, along with more severe functional impairment post-hospital discharge [15–17]. The Japanese Clinical Practice Guidelines for Management of Sepsis and Septic Shock 2020 (J-SSCG 2020) strongly emphasizes the importance of prevention and interventions to address PICS [18]. However, the evidence supporting these recommendations is derived from international studies, which involve different patient characteristics than those found in Japan, and therefore may not be applicable to the specific patient characteristics of critical care populations in Japan.

Consequently, we conducted a multicenter prospective cohort study involving 21 ICUs in tertiary hospitals

across Japan, aiming to investigate sepsis-related outcomes, including mortality and the incidence of PICS, for up to one year after hospital discharge.

Methods

Study settings

This was a multicenter prospective cohort study conducted across 21 ICUs in 20 tertiary hospitals in Japan. Ethical approvals were obtained from all hospitals (central ethics committee: Saiseikai Utsunomiya Hospital, Approval number: 2019–72). Informed consents were obtained from all patients. This study was registered in UMIN (UMIN000041433) and adhered to the STROBE guideline. The protocol for this study was published previously [19].

Patients

All consecutive patients admitted to the ICUs between 1 November 2020 and 30 April 2022 and who were diagnosed with sepsis or septic shock (according to the Sepsis-3 definition) at ICU admission were included. Exclusion criteria included patients who were under 18 years old, diagnosed with COVID-19, expected to be discharged from the ICU within 48 h, had a central nervous system disorder that was considered unrelated to sepsis based on clinical examination such as stroke, severe head trauma, brain tumor, hypoxic encephalopathy, cerebrovascular dementia, and Alzheimer's disease, could not communicate due to pre-existing psychiatric symptoms, could not walk independently even with a walking aid prior to hospitalization, or were in end-of-life or terminal state at ICU admission which could contribute to the treatment limitations in and after ICU stay. Patients for whom consent forms could not be obtained were also excluded. All patients at the participating ICUs received standard management based on the national sepsis guideline, (i.e., sepsis management, sedation, analgesia, delirium, etc.). [18, 20, 21].

Variables

The following data on patient characteristics were obtained at ICU admission, during the ICU stay, and at hospital discharge: age, sex, body mass index (BMI), Charlson Comorbidity Index (CCI) [22], Clinical Frailty Score (CFS) prior to hospital admission [23], Barthel Index [24] prior to hospital admission, recorded as the best value within the two weeks based on information from family members, employment status prior to hospital admission, ICU admission route, source of infection, Sequential Organ Failure Assessment (SOFA) sum score [25], presence of septic shock at ICU

admission, lactate level at ICU admission, and the use of organ support during the ICU stay, including noninvasive positive pressure ventilation (NPPV), high-flow nasal cannula (HFNC), invasive mechanical ventilation (IMV), and renal replacement therapy (RRT), ICU mortality, duration of IMV, and the length of ICU and hospital stay. Those who met the PICS Criteria, as described in the following section of *Outcome measures*, were also evaluated at the time of hospital discharge.

Outcome measures

Outcomes were assessed and obtained physically at hospital discharge and via telephone and mail at the 3-, 6-, and 12-month follow-ups after the hospital discharge by researchers from each hospital.

The primary outcome was a composite outcome as the rate of death (mortality) and the incidence of PICS, in consideration of the survival bias [26]. The incidence of PICS was defined by the presence of any of the following: (1) physical dysfunction, indicated by a score of 90 or lower on the Barthel Index (BI), (2) cognitive dysfunction, defined as a score of less than 40 on the Short Memory Questionnaire (SMQ) [27–29], or (3) a mental disorder, characterized by a score of 8 or higher on the anxiety or depression subscale of the Hospital Anxiety and Depression Scale (HADS) [30], or a score of 25 or higher on the Impact of Event Scale-Revised (IES-R) [31] at the follow-ups.

Among patients who survived to hospital discharge, (defined as survivors), the following secondary outcomes were obtained: the incidence of PICS, the incidence of each dysfunction, the score of each assessment, the number of patients with two of three or all three domains of PICS, EuroQoL 5-dimension 5-level (EQ-5D-5L) and visual analog scale (EQ-VAS) for the assessment of QOL [32], employment status, employment rates among those who were employed prior to hospitalization, body weight and changes compared to the baseline at ICU admission, readmissions to the hospital or ICU after discharge, unplanned emergency room visits, and the use of physical rehabilitation or psychiatric consultation clinics among survivors, as well as among those with physical dysfunctions or mental disorders, respectively, at each follow-up time point. The EQ-5D-5L was initially recorded as a 5-digit number and subsequently converted into the index value [27, 32, 33].

Statistical methods

Continuous variables were described using the median and interquartile range (IQR) and compared with the Mann–Whitney *U* test. Categorical variables were expressed as numbers and percentages and compared using the Fisher exact test or the Chi-square test as

appropriate. A multivariable logistic regression analysis was conducted to investigate the risk factors associated with each dysfunction at each follow-up time point. The covariates were selected as follows based on the clinically significant characteristics of the patients found in the current literature: age, sex (Male), BMI, CCI, CFS prior to hospital admission, BI prior to hospital admission, employment prior to hospital admission, ICU admission route of Emergency Room, source of Infection of abdomen, respiratory, urinary tract, musculoskeletal or soft tissue, SOFA sum score, septic shock, lactate level at the time of ICU admission, NPPV, HFNC, IMV, RRT, and length of hospital stay [29, 34–38]. In the multivariable analysis, the incidence of physical and cognitive dysfunction and mental disorders at the last follow-up were included as covariates. For instance, in the multivariable analysis examining the risk factors for physical dysfunction at the 12-month follow-up, the incidence of physical and cognitive dysfunction, along with mental disorders at the 6-month follow-up, was utilized as one of the covariates. To prevent overfitting in the multivariable analysis, we tested two additional models. The first model was analyzed using only the minimum variables, which included the incidence of physical and cognitive dysfunction, as well as mental disorders at the last follow-up. The second model included a small number of variables: Age, Sex, Body Mass Index, Clinical Frailty Scale, SOFA sum score, and the incidence of physical and cognitive dysfunction and mental disorders at the last follow-up. Missing data at each follow-up were not imputed in this analysis. All analyses were conducted using JMP software (version 13.0; SAS Institute Inc., Cary, NC, USA). Statistical tests were two-sided, and statistical significance was defined as a *p* value of less than 0.05.

Results

Patient characteristics

Out of 25,876 ICU admissions, 1382 patients were diagnosed with sepsis or septic shock at ICU admission during the study period, and a total of 339 patients were registered for the analysis of this study (Fig. 1).

The patients were old (median age of 74 years old [IQR: 67–82]), predominantly male (60%), and low BMI of 23.4 kg/m² [20.4–25.8]. They had few comorbidities (a median of 2 according to CCI), a median score of 3 in CFS [3–5], and a median of 100 in BI [90–100] prior to hospital admission. Around a quarter of patients (*n* = 80, 24%) were employed prior to hospital admission. The most common source of infection was the abdomen (32%), followed by the urinary tract (19%), musculoskeletal and soft tissue infections (19%), and respiratory infections (17%). At ICU admission, the SOFA score was a median




Fig. 1 Patient flow chart. COVID-19: Coronavirus disease 2019, ICU Intensive Care Unit

of 9 [6–12] and septic shock was identified in 77%. Additionally, 61% and 40% of the patients underwent IMV and RRT during ICU stay, respectively. The median duration of IMV was 5.0 days [2.8–10.0], with median length of ICU and hospital stays being 6.1 days [3.7–10.8], and 28.7 days [16.5–51.5], respectively.

Primary outcome

The incidence of death or those who met the PICS Criteria at hospital discharge was 89%, with the combined mortality and PICS incidence, the primary outcome, of 73%, 64%, and 65%, at the 3-, 6-, and 12-month follow-up, respectively (Fig. 2). Mortality consistently increased from 23% at hospital discharge to 37% at the 12-month follow-up. The proportion of patients who met the pre-defined PICS criteria at hospital discharge was 66%, reducing to a rate of 28% at the 12-month follow-up. The loss to follow-up rate during the period is shown in Supplemental Table 1. There was no difference in the characteristics of the patients among those who were followed up at each follow-up time point.

Outcomes among survivors

Among the survivors discharged from hospital, 85% met the criteria for PICS at the point of hospital discharge (Table 2). This reduced over the post-discharge period with reported rates of PICS as 62%,

47%, and 45% at the 3-, 6- and 12-month follow-ups. All three domains of PICS, including physical and cognitive dysfunction as well as mental disorders, consistently decreased during the first year of survivorship. Among the three symptoms of mental disorders, depression was consistently more prevalent over the year compared to anxiety or PTSD. The patients often experienced two or more dysfunctions simultaneously, decreasing from 59% of survivors at hospital discharge to 21% at the 12-month follow-up. The translated index value of EQ-5D-5L slightly improved over the year, while the EQ-VAS, a self-reported QOL value, showed less improvement. The overall employment status remained low over the 12 months following the discharge (15–17%), with slight reductions seen by 12 months in those survivors who had a job prior to hospitalization (52% at 3 months down to 44% at 12 months). Body weight was found to have decreased from baseline at the point of hospital discharge. Although gradual improvement was seen over the year following discharge from the hospital, body weight failed to return to baseline levels before the sepsis-related hospitalization. Two-fifths were readmitted to the hospital and one tenth were readmitted to ICU within the first year after discharge, and nearly one-third of the survivors utilized emergency services. The use of physical rehabilitation




Fig. 2 Primary outcome of the death or incidence of PICS. PICS Post Intensive Care Syndrome

or psychiatric consultation clinics remained consistently low during the first year, not only among all survivors but also among those who experienced physical dysfunction or mental disorders.

Risk factors

The incidence of physical, cognitive, or mental dysfunction was consistently and significantly associated with the incidence of the same dysfunction at the next follow-up (Table 3). This result remained consistent when the variables in the multivariable analysis were changed (Supplemental Table 2 and 3). Among the factors included in the multivariable analysis, age was consistently identified as an independent factor for physical and cognitive dysfunctions, and HFNC was for mental disorders (Supplemental Tables 4, 5, and 6).

Discussion

This multicenter prospective cohort study, conducted at 21 ICUs in 20 tertiary hospitals in Japan, demonstrated the characteristics of ICU patients with sepsis, the trajectories of their prognostic and functional outcomes (as a composite outcome of death or the incidence of PICS), and the impact of hospitalization on their post-sepsis lives over the first year after hospital discharge.

The results indicated that the patient characteristics were significantly different from those reported in previous studies of sepsis outside of Japan, highlighting the challenges of translating the past findings to clinical practice in Japan and underscoring the urgent need for studies conducted in Japan. Surprisingly, only one-third of ICU patients with sepsis survived without PICS, even a year after being discharged from the hospital. Furthermore, this situation did not improve between 6 and 12 months, indicating recovery from functional dysfunctions could reach a plateau within the first six months after hospital discharge.

The patients enrolled in this study were relatively older and had a lower BMI in comparison to previous sepsis studies, particularly those conducted in Western countries (Supplemental Table 7) [12, 13, 37, 39, 40]. Both advanced age and lower BMI independently correlate with long-term outcomes, including mortality and functional outcomes such as physical strength [41–45]. In addition, the patient cohort in this study was already on the brink of frailty based on the CFS value (a median of 3 [IQR: 3–5]) at ICU admission. This is likely due to the older age of our study population, which also serves as an independent factor affecting long-term outcomes [41, 46]. The source of infection was also different from sepsis-related studies conducted in Western countries, potentially due to the geographical or cultural background difference. These differences

Table 1 Characteristics of the patients

Variable	Overall (n = 339)
Baseline characteristics	
Age (year)	74 [67–82]
Sex (Male)	203 (60)
Body Mass Index (kg/m ²)	23.4 [20.4–25.8]
Charlson Comorbidity Index	2 [1–3]
Clinical Frailty Scale prior to hospital admission	3 [3–5]
Barthel Index prior to hospital admission	100 [90–100]
Employment status prior to hospital admission	80 (24)
ICU admission route	
ER	246 (73)
General ward	85 (25)
Others	8 (2)
Source of infection	
Abdomen	111 (32)
Urinary tract	63 (19)
Musculoskeletal and soft tissue	63 (19)
Respiratory	57 (17)
Other	45 (13)
Sequential Organ Failure Assessment (SOFA) sum score	9 [6–12]
Lactate level at ICU admission (mmol/L)	3.1 [1.6–5.8]
Presence of septic shock at ICU admission	261 (77)
Use of vasoactive drugs during ICU stay	
Noradrenaline	298 (88)
Vasopressin	145 (43)
Epinephrine	42 (12)
Use of medical devices during ICU stay	
Noninvasive positive pressure ventilation (NPPV)	27 (8)
High flow nasal cannula (HFNC)	57 (17)
Invasive mechanical ventilation (IMV)	208 (61)
Renal replacement therapy (RRT)	136 (40)
Clinical consequences of hospital stay	
ICU mortality	39 (12)
Duration of invasive mechanical ventilation	5.0 [2.8–10.0]
Length of ICU stay (days)	6.1 [3.7–10.8]
Length of Hospital stay (days)	28.7 [16.5–51.5]
Number of patients who met PICS criteria at hospital discharge	
Incidence of PICS	223 (66)
Incidence of physical dysfunction	164 (48)
Incidence of cognitive dysfunction	145 (43)
Incidence of mental disorder	129 (38)

The data in the table are presented as median with inter quartile range [IQR] or number with percentage (%)

ICU Intensive Care Unit, PICS Post-Intensive Care Syndrome

need to be taken into account when developing tailored interventions for the sepsis patient cohort in Japan [47]. Recent literature emphasizes that the variability and heterogeneity in patient cohorts highlight the necessity for personalized approaches that consider individual patient profiles, rather than adopting an one-size-fits-all

strategy [6, 36, 48]. This, however, can be also an alert when implementing evidence-based guidelines that were based on the findings from a patient cohort with distinct characteristics [49]. To promote a deeper understanding of long-term outcomes in the sepsis patient cohort in Japan and enhance their future care, a large-scale

Table 2 Outcomes among survivors

Item	Overall (n = 339)			
	Hospital discharge (n = 261)	3 month (n = 242)	6 month (n = 230)	12 month (n = 215)
Incidence of PICS among survivors	223 (85)	149 (62)	109 (47)	96 (45)
Incidence of physical dysfunction	164 (63)	79 (32)	55 (24)	42 (20)
A Score of BI	80 [40–100]	100 [75–100]	100 [85–100]	100 [90–100]
Incidence of cognitive dysfunction	145 (56)	85 (35)	57 (25)	63 (29)
A Score of SMQ	37 [27–42]	40 [33–45]	41 [36–45]	42 [36–45]
Incidence of mental disorder	129 (49)	77 (32)	68 (30)	50 (23)
Incidence of anxiety symptom	49 (19)	34 (14)	34 (15)	21 (10)
A score in the subscale for anxiety of HADS	4 [2–7]	4 [1–7]	3 [1–7]	2 [1–6]
Incidence of depression symptom	107 (41)	60 (25)	60 (26)	44 (20)
A score in the subscale for depression of HADS	7 [4–11]	5 [2–8]	5 [2–9]	5 [2–8]
Incidence of PTSD symptom	24 (9)	29 (12)	25 (11)	15 (7)
A Score of IES-R	4 [2–14]	7 [2–20]	7 [1–18]	6 [1–14]
Patients with two of three dysfunctions	81 (31)	39 (16)	25 (11)	30 (14)
Patients with all three domains of dysfunctions	72 (28)	27 (11)	23 (10)	15 (7)
EQ-5D-5L translated value	0.69 [0.42–0.83]	0.78 [0.55–0.89]	0.78 [0.60–0.89]	0.82 [0.68–1.00]
EQ-VAS	50 [7–70]	50 [10–80]	60 [8–80]	60 [30–80]
Employment status		36 (15)	34 (15)	37 (17)
Employment among those who had a job prior to the hospitalization ^a		34/66 (52)	30/66 (45)	27/62 (44)
Body weight (kg)		56 [47–64]	56 [49–65]	57 [48–66]
Changes in body weight as compared to the baseline (kg)		−4.0 [−0.6 ~ −8.9]	−3.2 [−8.1 ~ 0]	−2.0 [−6.0 ~ 1.4]
Changes in body weight as compared to the baseline (%)		−6.5 [−13.5 ~ −0.8]	−5.1 [−13.35 ~ 0]	−2.9 [−9.7 ~ 2.2]
Readmission to hospital after discharge		56 (24)	60 (28)	78 (41)
Readmission to ICU after discharge		8 (3)	12 (6)	16 (9)
Unplanned emergency room visits		19 (8)	37 (17)	58 (31)
Utilization of the physical rehabilitation clinic		20 (9)	30 (14)	28 (15)
Utilization of the physical rehabilitation clinic among those with physical dysfunction ^b		8/79 (10)	8/55 (15)	12/42 (29)
Utilization of the psychiatric consultation clinic		11 (5)	15 (7)	14 (7)
Utilization of the psychiatric consultation clinic among those with mental disorders ^c		9/77 (12)	8/68 (12)	7/50 (14)

Data in the table are presented as median with Inter Quartile Range (IQR) or number with percentage

BI Barthel Index, HADS Hospital Anxiety and Depression Scale, ICU Intensive Care Unit, IES-R Impact of Events Scale-Revised, EQ-5D-5L EuroQoL 5-dimension 5-level, PICS Post-Intensive Care Syndrome, SMQ Short Memory Questionnaire, VAS Visual Analogue Scale

^a Denominators represent the number of survivors who had a job prior to the hospitalization at the time of follow-ups

^b Denominators represent the number of survivors who had physical dysfunction at the time of follow-ups

^c Denominators represent the number of survivors who had mental disorders at the time of follow-ups

cohort registry with a comprehensive follow-up system is essential. In the future, Assistive Artificial Intelligence or robust machine learning methods might help identify patients at high risk of developing PICS and ultimately help to direct the prompt delivery of tailored interventions according to their risk assessment [50]. Currently, this study is ongoing to collect long-term outcomes for up to five years post-hospital discharge [19].

This study also revealed that a significant portion of ICU patients with sepsis either died or experienced serious functional deterioration which persisted for one year after hospital discharge. Only one-third of sepsis survivors were able to survive without PICS. Recovery appeared to plateau at six months after discharge, with no further significant change observed between six and twelve months, which is consistent with the recent report in the ICU population with COVID-19 infection in Japan

Table 3 Association of PICS at a follow-up with its prior follow-up

Variable	Physical dysfunction					
	3-month		6-month		12-month	
	Odds ratio	P value	Odds ratio	P value	Odds Ratio	P value
Incidence of physical dysfunction at the previous follow-up	7.21 [2.21–23.53]	< 0.01	24.24 [6.11–96.20]	< 0.01	21.41 [3.67–124.85]	< 0.01
Incidence of cognitive dysfunction at the previous follow-up	1.22 [0.41–3.65]	0.72	0.78 [0.20–3.04]	0.73	1.22 [0.16–9.07]	0.85
Incidence of mental disorder at the previous follow-up	1.42 [0.57–3.54]	0.46	3.15 [0.75–13.26]	0.12	1.12 [0.19–6.55]	0.90
Variable	Cognitive dysfunction					
	3-month		6-month		12-month	
	Odds ratio	P value	Odds ratio	P value	Odds Ratio	P value
Incidence of physical dysfunction at the previous follow-up	1.31 [0.53–3.23]	0.56	5.19 [1.31–20.61]	0.02	0.41 [0.10–1.65]	0.21
Incidence of cognitive dysfunction at the previous follow-up	3.97 [1.59–9.86]	< 0.01	16.29 [4.39–60.41]	< 0.01	4.12 [1.37–12.41]	0.01
Incidence of mental disorder at the previous follow-up	1.08 [0.48–2.41]	0.86	3.83 [1.05–14.02]	0.04	1.11 [0.37–3.30]	0.86
Variable	Mental disorders					
	3-month		6-month		12-month	
	Odds ratio	P value	Odds ratio	P value	Odds Ratio	P value
Incidence of physical dysfunction at the previous follow-up	2.29 [0.84–6.21]	0.11	2.09 [0.68–6.40]	0.20	0.95 [0.23–3.95]	0.95
Incidence of cognitive dysfunction at the previous follow-up	1.77 [0.69–4.55]	0.24	0.69 [0.25–1.88]	0.46	4.23 [1.24–14.44]	0.02
Incidence of mental disorder at the previous follow-up	4.30 [1.77–10.42]	< 0.01	10.37 [3.59–29.97]	< 0.01	7.18 [2.21–23.34]	< 0.01

The data in the table are presented as odds ratio with 95% confidence interval

[34]. Regarding the utilization of healthcare resources and services, a significant demand for resource utilization was evident, with a high rate of readmissions to the hospital or ICU, as well as visits to the emergency department. There was low utilization of healthcare services such as physical rehabilitation or psychiatric consultation clinics, even among those experiencing physical dysfunction or mental disorders. Given the substantially increasing burden on the healthcare system in terms of financial costs and the growing number of patients who continue to suffer after a critical illness [6], this imbalance urgently needs to be addressed through adequate support and appropriate resource allocation. As discussed that the overall incidence of PICS plateaued between 6 and 12 months, it is important to consider effective and prompt interventions soon or at least in the first 6 months after hospital discharge, not to miss the potentially critical period for recovery. Such actions will be crucial in facilitating survivors' return to their lives and reducing the healthcare burden. This is further supported by the finding that PICS-related dysfunctions independently contributed to the experience of PICS in the following three to six months. [6, 58, 59]. Investigating the effects of developed interventions in an appropriate manner, such as randomization, will be crucial to accelerate the flow toward their implementation in clinical settings. Future studies are also needed to verify whether developed

interventions can reduce healthcare resource usage and readmissions.

In this study, only small improvements were observed in QOL despite the almost halved incidence rate of PICS-associated dysfunctions among survivors. As shown in earlier studies, an improvement in symptoms associated with PICS did not lead to significant improvements in QOL scores (EQ-5D-5L) or self-reported QOL (EQ-VAS), suggesting that evaluating PICS-associated symptoms only is likely to be insufficient to assess their post-illness life [51, 52]. QOL could recover when other symptoms (i.e., PICS-associated dysfunctions) were first addressed, suggesting the last indicator to reflect overall recovery from sepsis. Therefore, QOL should be considered alongside assessments of PICS in future studies [53]. In this study, only half of those who were employed prior to hospitalization returned to work at three months, and around 20% of them lost their jobs between three and twelve months after discharge. This outcome was worse than earlier reports from Japan involving a general ICU patient cohort, where 20% of employed patients became unemployed by the twelve-month follow-up [54]. This may be due to sepsis resulting in a more challenging post-illness life than other ICU diseases because of its severity [13, 55]. Although our results do not clarify the relationship between job loss and PICS, this underscores the need for additional

support for ICU patients with sepsis to facilitate their return to and maintenance of employment [56, 57]. Further research, including qualitative interview surveys, would aid in understanding the key reasons behind job loss in the post-sepsis phase. Interestingly, the use of HFNC was associated with mental disorders. This may be, as previous reports suggested, because patients receiving HFNC were conscious during all medical events or procedures, which could be harmful, delusional, or lead to distorted memories, resulting in long-term health issues [60, 61].

Several limitations were acknowledged. First, the generalizability of this study requires attention when interpreting the results. This study involved a representative sample of 21 ICUs from 20 tertiary hospitals in Japan, covering only 7% of the ICU beds in the country. Given that the prevalence of sepsis in this study (5.3% of all ICU admissions) was similar to that in the previous cohort registry of ICU patients in Japan (4.3%) [14, 62], the findings of this study could have potential generalizability across Japan. Furthermore, the strength of this study lies in its exclusive focus on sepsis patients in the ICU, whereas the recent PICS study in Japan included a general ICU population, encompassing trauma and burn cases [28]. These cases tend to have a significantly different trajectory of functional recovery due to their specific disease nature and the procedures involved. The high follow-up rate should also be highlighted as the strength of this study, which could minimize the selection bias during the follow-up period. However, the exclusion of a large number of patients also needs to be taken into account. For example, the patients included in this study showed a low frequency of pulmonary infections, which was typically and frequently reported in the ICU population in Japan [63]. Second, the definition of PICS may vary between studies [27]. For example, while some prior studies have also employed the BI to assess physical function, others used different assessment tools. As a result, the incidence of PICS could differ when different assessment tools are applied. The results could also be biased by different assessors, especially when involving a large number of hospitals. Ideally, the outcome measurement should be performed by someone who is not involved in ICU care and not in a relationship with the patient or families to obtain objective outcomes. Nonetheless, the strength of this study lies in capturing a wide range of post-sepsis life, not only PICS-associated outcomes but also QOL, employment status after hospital discharge, and the utilization of healthcare resources and services. Third, the risk factors for PICS at each follow-up time point were identified in the multivariable analysis, although the causal relationship cannot be discussed within the constraints of this study's design. Furthermore, potential independent factors like prior psychiatric illness and social factors (i.e., family support status, economic conditions, etc.) were not collected

in this study. Fourth, this study was conducted during the COVID-19 pandemic, a time when hospitals and ICUs were significantly impacted. Thus, the current situation may differ from when this study was conducted, possibly resulting in a different post-sepsis life. To address these limitations, we propose a large-scale cohort registry of ICU sepsis in Japan with regular follow-up systems to comprehend the trajectories of post-sepsis life and develop effective interventions.

Conclusions

This multicenter prospective cohort study revealed the reality of post-sepsis recovery in ICU patients in Japan, who had a significantly different patient profile compared to previous PICS-related studies. Our results indicate major challenges exist in supporting the recovery of functions and improving the return to daily life in survivors of sepsis.

Abbreviations

ADL	Activities of daily living
BI	Barthel index
BMI	Body mass index
CCI	Charlson comorbidity index
CFS	Clinical frailty scale
COVID-19	Coronavirus disease 2019
EQ-5D-5L	EuroQoL 5-dimension 5-level
EQ-VAS	EuroQoL-visual analog scale
HADS	Hospital anxiety and depression scale
HFNC	High-flow nasal cannula
ICU	Intensive care unit
IES-R	Impact of events scale-revised
IMV	Invasive mechanical ventilation
IQR	Interquartile range
J-SSCG	Japanese clinical practice guidelines for management of sepsis and septic shock
NPPV	Noninvasive positive pressure ventilation
PICS	Post-intensive care syndrome
QOL	Quality of Life
RRT	Renal replacement therapy
SMQ	Short memory questionnaire
SOFA	Sequential organ failure assessment
STROBE	Strengthening the reporting of observational studies in epidemiology
UMIN	University hospital medical information network

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s40560-025-00792-0>.

Additional file 1

Additional file 2. Table 1: Follow-up loss rate at each time point. Table 2. Association of PICS at a follow-up with its prior follow-up in the multivariable analysis with the minimum variable. Table 3. Association of PICS at a follow-up with its prior follow-up in the multivariable analysis with a few variables. Table 4. Risk factors for the incidence of physical dysfunction at the time of each follow-up. Table 5. Risk factors for the incidence of cognitive dysfunction at the time of each follow-up. Table 6. Risk factors for the incidence of mental disorders at the time of each follow-up. Table 7. Differences in patient characteristics from past papers

Acknowledgements

We wish to thank the Japanese Association for Acute Medicine. We also thank the members of the ILOSS Study Committee Office: Motoki Sato, Rieko Kawauchi, Akiko Senba, Makoto Saita, and Hiromu Tsurumi who helped this project to be conducted smoothly. We thank all the non-author collaborators of the ILOSS Study Team listed in the Appendix "ILOSS_Study_Team-2024-03-30".

Author contributions

Study conception and design: KL, KN, HK, TK, and TO. Statistical analysis, or interpretation of data: KL, TK, DM, PN, SJS, and TG. Drafting the manuscript: KL, SW, KN, HN, MM, HK MA, KI, YM, TH, NS, YT, MH, TH, KO, DK, YS, YI, MN, KA, DS, HK, TK, DM, PN, SJS, and TO. Critical review and revision of the manuscript for important intellectual insight: SW, KN, HN, MM, HK MA, KI, YM, TH, NS, YT, MH, TH, KO, DK, YS, YI, MN, KA, DS, HK, TK, DM, PN, SJS, and TO. Study supervision: TK, PN, SJS, and TO. Edits and Reviews by a native English speaker: DM. All authors drafted the manuscript for important intellectual content, contributed to the revision of the final version of the manuscript, approved the final version submitted, and agreed to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved. TK and TO are the guarantors of the study. The corresponding author confirmed that all authors meet authorship criteria according to the International Committee of Medical Journal Editors (ICMJE).

Funding

This project is supported by the Japanese Association for Acute Medicine.

Availability of data and materials

No datasets were generated or analysed during the current study.

Declarations

Ethics approval and consent to participate

This study received ethics approvals from Saiseikai Utsunomiya Hospital (2020–42) and all other participating institutions. It was registered in the University Hospital Medical Information Network Clinical Trials Registry (Trial registration number: UMIN 000041433), and informed consent was obtained from all patients.

Consent for publication

Not applicable.

Competing interests

The authors declare no competing interests.

Author details

¹Non-Profit Organization ICU Collaboration Network (ICON), Tokyo, Japan. ²Department of Physical Therapy, Gifu University of Health Science, Gifu, Japan. ³Department of Rehabilitation, National Hospital Organization Nagoya Medical Center, Nagoya, Aichi, Japan. ⁴Department of Critical Care Medicine, Yokohama City University Hospital, 3-9, Fukuura, Kanazawa-Ku, Yokohama, Kanagawa 236-0004, Japan. ⁵Department of Emergency and Critical Care Medicine, Hitachi General Hospital, Hitachi, Ibaraki, Japan. ⁶Department of Emergency and Critical Care Medicine, Shinshu University School of Medicine, Nagano, Japan. ⁷Department of Emergency and Critical Care Medicine Faculty, Saga University Hospital, Saga, Japan. ⁸Department of Anesthesiology, Intensive Care Unit, Fukuyama City Hospital, Fukuyama, Hiroshima, Japan. ⁹Department of Emergency and Intensive Care Medicine, National Hospital Organization Nagoya Medical Center, Nagoya, Aichi, Japan. ¹⁰Department of Emergency, Critical Care, and Disaster Medicine, Okayama University Graduate School of Medicine, Dentistry, and Pharmaceutical Sciences, 2-5-1 Shikata-Cho, Okayama Kita-Ku, Okayama 700-8558, Japan. ¹¹Department of Emergency, Okayama Saiseikai General Hospital, 2-25 Kokutaityo, Okayama Kita-Ku, Okayama 700-8511, Japan. ¹²Emergency and Critical Care Medicine, Faculty of Medicine, University of Tsukuba, Ibaraki, Japan. ¹³Department of Emergency, Tsukuba Medical Center Hospital, Tsukuba, Ibaraki, Japan. ¹⁴Department of Rehabilitation, Japan Red Cross Narita Hospital, Narita, Chiba, Japan. ¹⁵Tajima Emergency & Critical

Care Medical Center, Toyooka Public Hospital, Toyooka, Hyogo, Japan.

¹⁶Department of Rehabilitation, Tsuchiura Kyodo General Hospital, Tsuchiura, Ibaraki, Japan. ¹⁷Department of Emergency and Critical Care Medicine, Nagoya University Graduate School of Medicine, Nagoya, Aichi, Japan. ¹⁸Department of Intensive Care Medicine, Okinawa Kyodo Hospital, Naha, Okinawa, Japan. ¹⁹Department of Emergency and Critical Care Medicine, Fukuoka University Hospital, Fukuoka, Fukuoka, Japan. ²⁰Department of Intensive Care Unit, Niigata University Medical and Dental Hospital, Niigata, Japan. ²¹Department of Emergency Medicine, SUBARU Health Insurance Society Ota Memorial Hospital, Ota, Gunma, Japan. ²²Graduate School of Data Science, Shiga University, Shiga, Japan. ²³Japanese Society for Early Mobilization, Tokyo, Japan. ²⁴Showa Medical University, Shinagawa, Tokyo, Japan. ²⁵Centre for Care Excellence, Coventry University, Coventry, UK. ²⁶Critical Care, University Hospitals Coventry & Warwickshire NHS Trust, Coventry, UK. ²⁷Nursing Research, University Hospital Schleswig-Holstein, Kiel, Germany. ²⁸Institute of Nursing Science and Development, Paracelsus Medical University, Salzburg, Austria. ²⁹Department of Anesthesia, Intensive Care Medicine and Pain Medicine, Clinical Division of General Anesthesia and Intensive Care Medicine, Medical University of Vienna, Wien, Austria. ³⁰Department of Anesthesiology and Intensive Care Medicine (CCM/CKV), Charité - Universitätsmedizin, Berlin, Germany. ³¹Department of Emergency Medicine and Critical Care Medicine, Tochigi Prefectural Emergency and Critical Care Center, Saiseikai Utsunomiya Hospital, Utsunomiya, Tochigi, Japan. ³²Present Address: 2-15-13 Hongo, Bunkyo-Ku, Tokyo 113-0033, Japan.

Received: 25 February 2025 Accepted: 17 April 2025

Published online: 01 May 2025

References

- Angus DC, van der Poll T. Severe sepsis and septic shock. *N Engl J Med*. 2013;369:840–51.
- Meyer NJ, Prescott HC. Sepsis and septic shock. *N Engl J Med*. 2024;391:2133–46.
- Rudd KE, Johnson SC, Agesa KM, Shackelford KA, Tsoi D, Kievlan DR, et al. Global, regional, and national sepsis incidence and mortality, 1990–2017: analysis for the global burden of disease study. *Lancet*. 2020;395:200–11.
- Kaukonen K-M, Bailey M, Suzuki S, Pilcher D, Bellomo R. Mortality related to severe sepsis and septic shock among critically ill patients in Australia and New Zealand, 2000–2012. *JAMA*. 2014;311:1308–16.
- Shankar-Hari M, Rubenfeld GD. Understanding long-term outcomes following sepsis: implications and challenges. *Curr Infect Dis Rep*. 2016;18:37.
- Inoue S, Nakanishi N, Amaya F, Fujinami Y, Hatakeyama J, Hifumi T, et al. Post-intensive care syndrome: recent advances and future directions. *Acute Med Surg*. 2024;11:e929.
- Harvey MA, Davidson JE. Postintensive care syndrome: right care, right now...and later. *Crit Care Med*. 2016;44:381–5.
- Desai SV, Law TJ, Needham DM. Long-term complications of critical care. *Crit Care Med*. 2011;39:371–9.
- Cuthbertson BH, Elders A, Hall S, Taylor J, MacLennan G, Mackirdy F, et al. Mortality and quality of life in the five years after severe sepsis. *Crit Care*. 2013;17:R70.
- Kurematsu K, Ikematsu Y. Quality of life of sepsis survivors. *Crit Care Med*. 2023;51:1339–49.
- Rhee C, Klompas M. Sepsis trends: increasing incidence and decreasing mortality, or changing denominator? *J Thorac Dis*. 2020;12:S89–100.
- Fleischmann-Struzek C, Born S, Kesselmeier M, Ely EW, Töpfer K, Romeike H, et al. Functional dependence following intensive care unit-treated sepsis: three-year follow-up results from the prospective Mid-German Sepsis Cohort (MSC). *Lancet Reg Health Eur*. 2024;46: 101066.
- Hodgson CL, Higgins AM, Bailey M, Barrett J, Bellomo R, Cooper DJ, et al. Comparison of 6-month outcomes of sepsis versus non-sepsis critically ill patients receiving mechanical ventilation. *Crit Care*. 2022;26:174.
- Imaeda T, Nakada T-A, Takahashi N, Yamao Y, Nakagawa S, Ogura H, et al. Trends in the incidence and outcome of sepsis using data from a Japanese nationwide medical claims database—the Japan Sepsis Alliance (JaSA) study group. *Crit Care*. 2021;25:338.

15. Bruno RR, Wernly B, Mamandipoor B, Rezar R, Binnebössel S, Baldia PH, et al. ICU-mortality in old and very old patients suffering from sepsis and septic shock. *Front Med (Lausanne)*. 2021;8: 697884.
16. Rowe T, Araujo KLB, Van Ness PH, Pisani MA, Juthani-Mehta M. Outcomes of older adults with sepsis at admission to an intensive care unit. *Open Forum Infect Dis*. 2016;3:ofw010.
17. Kotfis K, Wittebole X, Jaschinski U, Solé-Violán J, Kashyap R, Leone M, et al. A worldwide perspective of sepsis epidemiology and survival according to age: observational data from the ICON audit. *J Crit Care*. 2019;51:122–32.
18. Egi M, Ogura H, Yatabe T, Atagi K, Inoue S, Iba T, et al. The Japanese clinical practice guidelines for management of sepsis and septic shock 2020 (J-SSCG 2020). *J Intensiv Care Med*. 2021;9:53.
19. Liu K, Kotani T, Nakamura K, Chihiro T, Morita Y, Ishii K, et al. Effects of evidence-based ICU care on long-term outcomes of patients with sepsis or septic shock (ILOSS): protocol for a multicentre prospective observational cohort study in Japan. *BMJ Open*. 2022;12: e054478.
20. Rhodes A, Evans LE, Alhazzani W, Levy MM, Antonelli M, Ferrer R, et al. Surviving sepsis campaign: international guidelines for management of sepsis and septic shock: 2016. *Intensive Care Med*. 2017;43:304–77.
21. Devlin JW, Skrobik Y, Gélinas C, Needham DM, Slooter AJC, Pandharipande PP, et al. Clinical practice guidelines for the prevention and management of pain, agitation/sedation, delirium, immobility, and sleep disruption in adult patients in the ICU. *Crit Care Med*. 2018;46:e825–73.
22. Charlson ME, Carrozzino D, Guidi J, Patierno C. Charlson comorbidity index: a critical review of clinimetric properties. *Psychother Psychosom*. 2022;91:8–35.
23. Church S, Rogers E, Rockwood K, Theou O. A scoping review of the Clinical Frailty Scale. *BMC Geriatr*. 2020;20:393.
24. Mahoney FI, Barthel DW. Functional evaluation: the Barthel index. *Md State Med J*. 1965;14:61–5.
25. Vincent JL, Moreno R, Takala J, Willatts S, De Mendonça A, Bruining H, et al. The SOFA (sepsis-related organ failure assessment) score to describe organ dysfunction/failure on behalf of the working group on sepsis-related problems of the European society of intensive care medicine. *Intensiv Care Med*. 1996;22:707–10.
26. Hodgson CL, Higgins AM, Bailey MJ, Anderson S, Bernard S, Fulcher BJ, et al. Incidence of death or disability at 6 months after extracorporeal membrane oxygenation in Australia: a prospective, multicentre, registry-embedded cohort study. *Lancet Respir Med*. 2022;10:1038–48.
27. Nakanishi N, Liu K, Kawauchi A, Okamura M, Tanaka K, Katayama S, et al. Instruments to assess post-intensive care syndrome assessment: a scoping review and modified Delphi method study. *Crit Care*. 2023;27:430.
28. Kawakami D, Fujitani S, Morimoto T, Dote H, Takita M, Takaba A, et al. Prevalence of post-intensive care syndrome among Japanese intensive care unit patients: a prospective, multicenter, observational J-PICS study. *Crit Care*. 2021;25:69.
29. Hatakeyama J, Inoue S, Liu K, Yamakawa K, Nishida T, Ohshimo S, et al. Prevalence and risk factor analysis of post-intensive care syndrome in patients with COVID-19 requiring mechanical ventilation: a multicenter prospective observational study. *J Clin Med Res*. 2022;11:5758.
30. Zigmond AS, Snaith RP. The hospital anxiety and depression scale. *Acta Psychiatr Scand*. 1983;67:361–70.
31. Asukai N, Kato H, Kawamura N, Kim Y, Yamamoto K, Kishimoto J, et al. Reliability and validity of the Japanese-language version of the impact of event scale-revised (IES-R-J): four studies of different traumatic events. *J Nerv Ment Dis*. 2002;190:175–82.
32. Shiroya T, Fukuda T, Ikeda S, Igarashi A, Noto S, Saito S, et al. Japanese population norms for preference-based measures: EQ-5D-3L, EQ-5D-5L, and SF-6D. *Qual Life Res*. 2016;25:707–19.
33. Tanaka S, Ando K, Kobayashi K, Seki T, Hamada T, Machino M, et al. Association between locomotive syndrome and the Japanese version of the EQ-5D-5L in middle-aged and elderly people in Japan. *Nagoya J Med Sci*. 2020;82:5–14.
34. Hatakeyama J, Nakamura K, Inoue S, Liu K, Yamakawa K, Nishida T, et al. Two-year trajectory of functional recovery and quality of life in post-intensive care syndrome: a multicenter prospective observational study on mechanically ventilated patients with coronavirus disease-19. *J Intensiv Care*. 2025;13:7.
35. Liu K, Ogura T, Takahashi K, Nakamura M, Ohtake H, Fujiduka K, et al. A Progressive early mobilization program is significantly associated with clinical and economic improvement: a single-center quality comparison study. *Crit Care Med*. 2019;47:e744–52.
36. Liu K, Tronstad O, Flaws D, Churchill L, Jones AYM, Nakamura K, et al. From bedside to recovery: exercise therapy for prevention of post-intensive care syndrome. *J Intensiv Care Med*. 2024;12:11.
37. Mesina RS Jr, Rustøen T, Hagen M, Laake JH, Hofso K. Long-term functional disabilities in intensive care unit survivors: a prospective cohort study. *Aust Crit Care*. 2024;37:843–50.
38. Watanabe S, Liu K, Nakamura K, Koza R, Horibe T, Ishii K, et al. Association between early mobilization in the ICU and psychiatric symptoms after surviving a critical illness: a multi-center prospective cohort study. *J Clin Med Res*. 2022;11:2587.
39. Maley JH, Sandsmark DK, Trainor A, Bass GD, Dabrowski CL, Magdamo BA, et al. Six-month impairment in cognition, mental health, and physical function following COVID-19-associated respiratory failure. *Crit Care Explor*. 2022;4: e0673.
40. Liu K, Nakashima T, Goto T, Nakamura K, Nakano H, Motoki M, et al. Phenotypes of functional decline or recovery in sepsis ICU survivors: Insights from a 1-year follow-up multicenter cohort analysis. *Crit Care Med*. 2025. <https://doi.org/10.1097/CCM.0000000000006621>.
41. Watanabe D, Yoshida T, Watanabe Y, Yamada Y, Kimura M. Is a higher body mass index associated with longer duration of survival with disability in frail than in non-frail older adults? *Int J Obes (Lond)*. 2025;49:348–56.
42. Winter JE, MacInnis RJ, Wattanapenpaiboon N, Nowson CA. BMI and all-cause mortality in older adults: a meta-analysis. *Am J Clin Nutr*. 2014;99:875–90.
43. Woolley C, Thompson C, Hakendorf P, Horwood C. The effect of age upon the interrelationship of BMI and inpatient health outcomes. *J Nutr Health Aging*. 2019;23:558–63.
44. Martin GS, Mannino DM, Moss M. The effect of age on the development and outcome of adult sepsis. *Crit Care Med*. 2006;34:15–21.
45. Martin-Loeches I, Guia MC, Vallecocchia MS, Suarez D, Ibarz M, Irazabal M, et al. Risk factors for mortality in elderly and very elderly critically ill patients with sepsis: a prospective, observational, multicenter cohort study. *Ann Intensive Care*. 2019;9:26.
46. Wozniak H, Beckmann TS, Dos Santos RA, Pugin J, Heidegger C-P, Cereghetti S. Long-stay ICU patients with frailty: mortality and recovery outcomes at 6 months. *Ann Intensive Care*. 2024;14:31.
47. Sato T, Kudo D, Kushimoto S, Hasegawa M, Ito F, Yamanouchi S, et al. Associations between low body mass index and mortality in patients with sepsis: A retrospective analysis of a cohort study in Japan. *PLoS ONE*. 2021;16: e0252955.
48. Wang RC, Wang Z. Precision medicine: disease subtyping and tailored treatment. *Cancers*. 2023;15:3837.
49. Costa DK, White MR, Ginier E, Manojlovich M, Govindan S, Iwashyna TJ, et al. Identifying barriers to delivering the awakening and breathing coordination, delirium, and early exercise/mobility bundle to minimize adverse outcomes for mechanically ventilated patients: a systematic review. *Chest*. 2017;152:304–11.
50. Alowais SA, Alghamdi SS, Alsuehaby N, Alqahtani T, Alshaya AI, Almohareb SN, et al. Revolutionizing healthcare: the role of artificial intelligence in clinical practice. *BMC Med Educ*. 2023;23:689.
51. Nannan Panday RS, Minderhoud TC, Chantalou DS, Alam N, Nanayakkara PWB. Health related quality of life in sepsis survivors from the prehospital antibiotics against sepsis (PHANTASI) trial. *PLoS ONE*. 2019;14: e0222450.
52. Alam N, Nannan Panday RS, Heijnen JR, van Galen LS, Kramer MH, Nanayakkara PWB. Long-term health related quality of life in patients with sepsis after intensive care stay: a systematic review. *Acute Med*. 2017;16: 164–9.
53. Kjær M-BN, Bruun CRL, Granholm A, Møller MH, Rasmussen BS, Mortensen CB, et al. A core outcome set for adult general ICU patients. *Crit Care Med*. 2025;53:e575–89.
54. Unoki T, Kitayama M, Sakuramoto H, Ouchi A, Kuribara T, Yamaguchi T, et al. Employment status and its associated factors for patients 12 months after intensive care: Secondary analysis of the SMAP-HoPe study. *PLoS ONE*. 2022;17: e0263441.
55. Sheu C-C, Gong MN, Zhai R, Chen F, Bajwa EK, Clardy PF, et al. Clinical characteristics and outcomes of sepsis-related vs non-sepsis-related ARDS. *Chest*. 2010;138:559–67.

56. Hatakeyama J, Nakamura K, Sumita H, Kawakami D, Nakanishi N, Kashiwagi S, et al. Intensive care unit follow-up clinic activities: a scoping review. *J Anesth*. 2024;38:542–55.
57. Nakanishi N, Liu K, Hatakeyama J, Kawauchi A, Yoshida M, Sumita H, et al. Post-intensive care syndrome follow-up system after hospital discharge: a narrative review. *J Intensiv Care Med*. 2024;12:2.
58. Shirasaki K, Hifumi T, Nakanishi N, Nosaka N, Miyamoto K, Komachi MH, et al. Postintensive care syndrome family: a comprehensive review. *Acute Med Surg*. 2024;11: e939.
59. Inoue S, Hatakeyama J, Kondo Y, Hifumi T, Sakuramoto H, Kawasaki T, et al. Post-intensive care syndrome: its pathophysiology, prevention, and future directions. *Acute Med Surg*. 2019;6:233–46.
60. Danielis M, Movio F, Milanese G, Mattiussi E. Patients' reports on their delusional memories from the intensive care unit: a systematic review of qualitative studies. *Intensiv Crit Care Nurs*. 2024;81: 103617.
61. Fukuda T, Watanabe N, Sakaki K, Monna Y, Terachi S, Miyazaki S, et al. Identifying cues of distorted memories in intensive care by focus group interview of nurses. *Nurs Open*. 2022;9:666–75.
62. Imaeda T, Oami T, Takahashi N, Saito D, Higashi A, Nakada T-A. Epidemiology of sepsis in a Japanese administrative database. *Acute Med Surg*. 2023;10: e890.
63. Ogura H, Gando S, Saitoh D, Takeyama N, Kushimoto S, Fujishima S, et al. Epidemiology of severe sepsis in Japanese intensive care units: a prospective multicenter study. *J Infect Chemother*. 2014;20:157–62.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.